



EFRE Recurrent Implantation Failure (RIF) Survey


Please Scan the QR Code






1- In your practice, you define RIF according to?

24 responses

 Copy








-  A- Number of previously failed fresh cycles
-  B- Number of previously failed fresh and frozen cycles
-  C- Number of embryos previously failed to implant

2- In your practice, you define RIF in cases with

24 responses

 Copy




-  A- Previously failed 1 IVF/ICSI cycle.
-  B- Previously failed 2 IVF/ICSI cycle.
-  C- Previously failed 3 IVF/ICSI cycle.
-  D- Previously failed 4 IVF/ICSI cycle.

3- Do you take the female age in consideration in the definition of RIF?

 Copy


24 responses





4- Do you take the quality of embryos transferred in consideration in the definition of RIF?

 Copy

24 responses




 A- YES
 B- NO

5- In your practice, how do you assess the uterine cavity in RIF cases?



24 responses








- A- 2 D ultrasound.
- B- 3 D ultrasound.
- C- Office hysteroscopy (without sedation nor anesthesia).
- D- Diagnostic hysteroscopy (with sedation or anesthesia).

6- When do you order thrombophilia screening?

 Copy

24 responses




-  A- Before the first ICSI trial.
-  B- After 1 failed ICSI trial.
-  C- After 2 failed ICSI trial.
-  D- After 3 failed ICSI trial.

7- When do you order karyotyping (male/female)?

 Copy

24 responses









- A- Never
- B- Before the first ICSI trial.
- C- After 1 failed ICSI trial.
- D- After 2 failed ICSI trial.
- E- After 3 failed ICSI trial.

8- When do you order serum vit D?

 Copy

24 responses




-  A- Never
-  B- Before the first ICSI trial.
-  C- After 1 failed ICSI trial.
-  D- After 2 failed ICSI trial.
-  E- After 3 failed ICSI trial.

9- When do you order sperm DNA fragmentation?

 Copy

24 responses








- A- Never
- B- Before the first ICSI trial.
- C- After 1 failed ICSI trial.
- D- After 2 failed ICSI trial.
- E- After 3 failed ICSI trial.

10- In my practice, I prescribe low dose aspirin for:

 Copy

24 responses








-  A- All ICSI cases.
-  B- All RIF cases
-  C- Cases with abnormal thrombophilia screening
-  D- Never

11- In my practice, I prescribe low molecular weight heparin for:

 Copy

24 responses







-  A- All ICSI cases.
-  B- All RIF cases
-  C- Cases with abnormal thrombophilia screening
-  D- Never

12- In my practice, I prescribe corticosteroids for:

 Copy

24 responses



-  A- All ICSI cases.
-  B- RIF cases
-  C- Never

13- Which of the following procedures you perform for RIF cases? (you can choose more than one)



24 responses

